# Genome-wide association study identifies *BTNL2* associated with atopic asthma in children

Soo Yeon Kim, MD<sup>a</sup>, Eun Gyul Kim, MS<sup>a</sup>, Mina Kim, PhD<sup>a</sup>, Jung Yeon Hong, PhD<sup>a,b</sup>, Ga Eun Kim, MD<sup>a,c</sup>, Jae Hwa Jung, MD<sup>a</sup>, Mireu Park, MD<sup>a</sup>, Min Jung Kim, MD, PhD<sup>a</sup>, Yoon Hee Kim, MD, PhD<sup>a</sup>, Myung Hyun Sohn, MD, PhD<sup>a</sup>, Kyung Won Kim, MD, PhD<sup>a,\*</sup>

## Abstract

Asthma is a heterogeneous disease characterized by chronic airway inflammation with a genetic predisposition. Butyrophilin-like 2 (*BTNL2*) is a member of the immunoglobulin superfamily that plays an important role in regulating T cell activation and immune homeostasis. Here, we aimed to investigate the association of the genetic variants of *BTNL2* with childhood asthma and asthmarelated traits by utilizing extreme asthma phenotypes and employing a genome-wide association study. Our study included 243 children with well-defined moderate to severe atopic asthma and 134 healthy children with no history of allergic diseases and allergic sensitization. DNA from these subjects was genotyped using Axiom<sup>TM</sup> Genome-Wide Array Plates. Although no single nucleotide polymorphisms (SNPs) reached a genome-wide threshold of significance, 3 SNPs, rs3817971, rs41355746, and rs41441651, at *BTNL2* were significantly associated with moderate to severe atopic asthma after performing Bonferroni correction. These SNPs were also associated with the risk of allergic sensitization toward house dust mites and the presence and degree of bronchial hyperresponsiveness. Thus, we identified that *BTNL2* was associated with atopic moderate to severe persistent asthma in Korean children, and this may play an important role in disease development and susceptibility.

**Abbreviations:** BDR = bronchodilator response, BHR = bronchial hyperresponsiveness, BTNL2 = Butyrophilin-like 2, Der f = *Dermatophagoides farina*, Der p = *Dermatophagoides pteryonyssinus*, FEV<sub>1</sub> = forced expiratory volume in 1 s, GWAS = genomewide association study, IgE = immunoglobulin E, LD = linkage disequilibrium, SNP = single nucleotide polymorphism.

Keywords: allergic sensitization, asthma, bronchial hyperresponsiveness, children, genome-wide association study

Editor: Sabahat Sarfaraz.

This work was supported by a faculty research grant of Yonsei University College of Medicine (6-2012-0016) and the National Research Foundation Grant funded by the Korean Government (NRF-2019R1F1A1058910).

The study was conducted according to the guidelines of the Declaration of Helsinki, and approved by the Institutional Review Board of SEVERANCE HOSPITAL (IRB no. 4-2004-0036).

Informed consent was obtained from all subjects involved in the study.

The authors declare no conflict of interest.

The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

Supplemental Digital Content is available for this article.

<sup>a</sup> Department of Pediatrics, Severance Hospital, Institute of Allergy, Brain Korea 21 Project for Medical Science, Yonsei University College of Medicine, Seoul, Republic of Korea, <sup>b</sup> Division of Cardiovascular Disease Research, Department for Chronic Disease Convergence Research, Korea National Institute of Health, Cheongju, Republic of Korea, <sup>c</sup> Department of Pediatrics, Keimyung University School of Medicine, Keimyung University Dongsan Hospital, Daegu, Republic of Korea.

\* Correspondence: Kyung Won Kim, Department of Pediatrics, Institute of Allergy, Severance Children's Hospital, Yonsei University College of Medicine, 50-1 Yonsei-ro, Seodaemun-gu, Seoul 03722, Korea (e-mail: kwkim@yuhs.ac).

Copyright © 2021 the Author(s). Published by Wolters Kluwer Health, Inc. This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial License 4.0 (CCBY-NC), where it is permissible to download, share, remix, transform, and buildup the work provided it is properly cited. The work cannot be used commercially without permission from the journal.

How to cite this article: Kim SY, Kim EG, Kim MN, Hong JY, Kim GE, Jung JH, Park M, Kim MJ, Kim YH, Sohn MH, Kim KW. Genome-wide association study identifies BTNL2 associated with atopic asthma in children. Medicine 2021;100:44(e27626).

Received: 18 April 2021 / Received in final form: 18 August 2021 / Accepted: 6 October 2021

http://dx.doi.org/10.1097/MD.000000000027626

## 1. Introduction

Asthma is a heterogenous and genetically complex respiratory disease that is a result of gene–environment interactions, especially early in life.<sup>[1]</sup> Gene discovery approaches in asthma were initially knowledge-based candidate gene association studies and family-based genome-wide linkage analyses. Recently, genome-wide association studies (GWASs) have dominated; these studies aid hypothesis-free discovery of novel risk loci, which in turn provide mechanistic insights into disease pathogenesis.<sup>[2–4]</sup>

Medicine

GWASs of asthma have identified many risk loci, including 17q12-21 (*ORMDL3*, *GSDMB*), 6p21 (HLA region), 2q12 (*IL1RL1/IL18R1*), 5q22 (*TSLP*), and 9p24 (*IL33*) that provided novel insights into asthma biology.<sup>[4]</sup> In addition, GWASs have revealed significant genome-wide variants for asthma-related traits, such as bronchodilator response (BDR),<sup>[5]</sup> bronchial hyperresponsiveness (BHR),<sup>[6]</sup> blood eosinophils,<sup>[7]</sup> total serum immunoglobulin E (IgE) levels,<sup>[2,8]</sup> and allergic sensitizations.<sup>[9]</sup> Such studies are based on the assumptions that it is easy to detect genes influencing components of asthma by reducing asthma heterogeneity and that the same genes also contribute to asthma risk.

However, most previous GWASs of asthma comprise subjects of European ancestry; only a few studies have been conducted in ethnic minority populations. As populations vary with respect to allele frequencies, linkage disequilibrium (LD) patterns, and effect sizes of variants underlying disease risk,<sup>[10–12]</sup> the previous GWASs may have limited utility in other population groups. Furthermore, they have probably missed important risk variants in non-European populations.

In the Korean population, GWASs of asthma have focused on occupation-associated forms of asthma and aspirin-exacerbated

respiratory disease in adults.<sup>[13–17]</sup> A recent GWAS in Korean subjects with asthma–chronic obstructive pulmonary disease overlap syndrome did not reveal any significant genome-wide hits.<sup>[18]</sup> Additionally, no significant genome-wide loci were reported in a GWAS based on total serum IgE levels in Korean subjects with asthma.<sup>[19]</sup> Such limited GWAS results in the Korean population are most likely because of the small sample size. Considering that GWAS adopts rigorous statistical control for false-discovery rate, which may miss several single nucleotide polymorphisms (SNPs) of true associations,<sup>[20]</sup> there is a need for alternative strategies to address such concerns.

Therefore, the aim of the present study was to identify asthmarelated loci using cases of extreme phenotypes – that is, moderateto-severe persistent atopic asthma – and healthy controls in a small-sized GWAS. In addition, we sought to determine the GWAS-based SNPs associated with asthma susceptibility traits, including BDR, BHR, and total or specific serum IgE levels.

#### 2. Methods

#### 2.1. Study subjects

For this case-control investigation, 243 subjects with moderate to severe persistent asthma with allergic sensitization, and 134 healthy children without a history of allergic disease or allergic sensitization were included in the study. All children were recruited from Severance Children's Hospital, Seoul, Korea during April, 2004 and May, 2010. The study was approved by the Institutional Review Board of Severance Hospital (IRB no. 4-2004-0036). All participants were unrelated, and either they or their parents provided written informed consent.

Subjects with asthma were confirmed based on consistent respiratory symptoms verified by physicians, presence of either a BDR of  $\geq 12\%$  increase in forced expiratory volume in 1 s (FEV<sub>1</sub>) or BHR of decrease in FEV<sub>1</sub> by  $\geq 20\%$  upon inhalation of <16 mg/mL methacholine. Allergic sensitization was defined by specific serum IgE levels greater than  $0.7 kU_A/L$  for at least one of the following food or airborne allergens: egg white, milk, *Dermatophagoides pteronyssinus* (Der p), *Dermatophagoides farina* (Der f), Alternaria species, or *Blattella germanica*. Children with moderate to severe asthma according to the Global Initiative for Asthma guidelines were recruited.

Healthy controls included children who had visited the hospital for a general health checkup or vaccination, and had no history of allergic diseases based on interviews with their parents. Further, children who were not sensitized to the abovementioned 6 common allergens and had total serum IgE levels <100 kU/L were enrolled.

The total and specific IgE levels to selected allergens were measured from peripheral blood samples using the ImmunoCAP system (Pharmacia Diagnostics, Uppsala, Sweden). Complete blood count, including eosinophil count, was analyzed using the ADVIA 2120i hematology system with autoslide (Siemens Healthcare Diagnostics Inc., Deerfield, IL, USA).

## 2.2. Genotyping, quality control, and Butyrophilin-like 2 (BTNL2) region selection

Blood samples were collected from each subject to extract genomic DNA. The DNA was then genotyped using an Affymetrix Axiom array (Affymetrix Inc., Santa Clara, CA, USA). There was originally a total of 600,252 SNPs; after frequency and genotyping pruning using GRCh37/hg19, there were 574,420 autosomal SNPs. During quality control filtration, samples with an autosomal SNP call rate <95% were removed. SNPs with minor allele frequencies <5% or Hardy–Weinberg equilibrium *P* value <10<sup>-6</sup> were also excluded. Finally, 423,461 SNPs with a genomic inflation factor ( $\lambda$ ) of 1.0033 were used for the GWAS (Supplemental Figure S1, http://links.lww.com/MD2/A616). Among the finally refined SNPs, we then selected 46 SNPs located ± 500 base pairs around *BTNL2* in order to further analyze the *BTNL2* region.

#### 2.3. Statistical analysis

Genome-wide associations between atopic asthma and variants were analyzed using PLINK (v.1.07).<sup>[21]</sup> Correlation coefficients were calculated and genome-wide statistical analyses were performed using PLINK. Genomic inflation factor ( $\lambda$ ) was calculated based on median chi-square statistics. Manhattan plots and quantile–quantile plots were plotted using the "manhattanly" and "qqman" libraries, respectively, in R (v.3.3.3).<sup>[22]</sup> LD calculations were performed using PLINK, and loci, including variants significantly associated with asthma, were plotted using LocusZoom (v.0.4.8).<sup>[23]</sup> Association of genetic polymorphisms with BHR and levels of total and mitespecific IgE were evaluated by logistic and linear regression analyses using the software R. Age and sex were the adjustment factors for the analyses.

#### 3. Results

#### 3.1. Subject characteristics

Demographic characteristics of the study subjects are summarized in Table 1. The control and asthma groups did not differ in

#### Table 1

#### Clinical profiles of the study subjects (N=377).

	Control	Atopic asthma	
	(n = 134)	(n=243)	P value
Sex, M (%)	63 (47.0)	168 (69.1)	<.001
Age (yr)	9.1 (7.3–11.6)	8.4 (6.5-10.5)	.051
Height (cm)	133.6 (125.0–147.0)	131.0 (119.2–142.0)	.034
Weight (kg)	31.0 (24.7-45.0)	30.0 (23.0-40.0)	.096
FVC (% pred)	92.8 (84.0-102.1)	91.9 (80.8-100.1)	.361
FEV <sub>1</sub> (% pred)	98.3±15.7	91.7±15.6	<.001
Post-BD FEV <sub>1</sub> (% pred)	108.2 ± 11.9	101.4±16.1	<.001
FEV <sub>1</sub> /FVC (%)	92.0 (87.3–97.3)	86.7 (79.3-93.2)	<.001
PEF (% pred)	94.5 (81.3-104.8)	88.0 (77.0-99.0)	.002
FEF <sub>25-75</sub> (% pred)	$101.9 \pm 23.6$	80.9±27.0	<.001
$\Delta FEV_1$	4.1 (1.1-6.1)	9.6 (4.2-15.3)	<.001
PC <sub>20</sub> (mg/mL)	N/A	3.7 (1.6-7.3)	N/A
PC <sub>20</sub> category			<.001
<1	0 (0.0)	42 (17.6)	
1-4	0 (0.0)	85 (35.6)	
4–16	0 (0.0)	103 (43.1)	
≧16	134 (100.0)	9 (3.8)	
Blood eosinophils (/µL)	140.0 (90.0-200.0)	520.0 (300.0-740.0)	<.001
Total serum IgE (kU/L)	28.9 (15.8-53.0)	584.0 (317.0-1035.0)	<.001

Data are given as number (%), mean ( $\pm$  standard deviation), or median (interquartile range).  $\Delta$ FEV<sub>1</sub> = change in FEV<sub>1</sub> after BD; BD = bronchodilator; FEF<sub>25-75</sub> = maximum mid-expiratory flow; FEV<sub>1</sub> = forced expiratory volume in 1 s; FVC = forced vital capacity; IgE = immunoglobulin E; N/A = not applicable; PC<sub>20</sub> = provocative concentration of methacholine inducing a 20% decrease in FEV<sub>1</sub>; PEF = peak expiratory flow.



Figure 1. Regional association plot of 6p21.3 locus for atopic moderate to severe persistent asthma. Statistical significance of each single nucleotide polymorphism (SNP) on the  $-\log_{10}$  scale (left *y*-axis) according to its chromosomal position (*x*-axis) is shown. Relative location of the genes in each region and the direction of transcription is shown in the bottom panel. The blue line shows recombination rate across that region (right y-axis). The most associated SNP (rs3817971) is shown as a purple circle, and the remaining SNPs are color-coded according to the level of linkage disequilibrium ( $r^2$ ).

age. Males were predominant in the asthma group (n=168, 69.1%). Blood eosinophil count and total serum IgE levels were significantly higher in the asthma group than in the control group (*P* value <.001 for both). Compared with healthy controls, children with asthma showed a lower percentage of FEV<sub>1</sub>/forced vital capacity ratio, percentage of predicted FEV<sub>1</sub>, percentage of predicted maximum mid-expiratory flow (FEF<sub>25-75</sub>), and peak expiratory flow (Table 1).

# 3.2. Genome-wide associations with moderate-to-severe persistent atopic asthma

A total of 423,461 autosomal SNPs that passed quality control were analyzed for their association with asthma. GWAS of atopic asthma and healthy control subjects showed an excess of small *P* values compared with those expected by chance (Supplemental Figure S1, http://links.lww.com/MD2/A616). Although no SNPs reached a genome-wide threshold of significance ( $P < 1.18 \times 10^{-7}$ ), several candidate loci were identified as top signals (Manhattan plot in Supplemental Figure S2, http://links.lww. com/MD2/A617); the most prominent SNP was located on chromosome 6p21.32. Figure 1 displays a regional association plot at the 6p21.32 locus for atopic asthma. The most highly associated SNP was rs3817971, which is located within *BTNL2*.

# 3.3. Association of genetic variations in BTNL2 with asthma

To analyze *BTNL2* polymorphisms, we selected SNPs that were located  $\pm 500$  base pairs around *BTNL2*. A total of 46 SNPs were reanalyzed, which yielded 3 SNPs (rs3817971, rs41355746, and rs41441651) with *P* values <.001087 (Bonferroni-corrected; Table 2 and Supplemental Table S1, http://links.lww.com/MD2/A618). Risk alleles of these SNPs were associated with an increased risk of asthma after age and sex adjustment (rs3817971: adjusted odds ratio [aOR]=3.314, *P*<.001; rs41355746: aOR=2.639, *P*=.001; rs41441651: aOR=2.404, *P*=.002). These associations between genetic polymorphisms and asthma remained significant for both additive and dominant models (Supplemental Table S2, http://links.lww.com/MD2/A619).

#### 3.4. Associations with BDR and BHR

A case-control analysis adjusted for age and sex as covariates was performed for determining association with additional phenotypes. Table 3 represents the association results of the provocative concentration of methacholine that caused a 20% drop in FEV<sub>1</sub> from baseline level (PC<sub>20</sub>). PC<sub>20</sub> levels were further Table 2

			Minor allel	e frequency			
SNP	Position	Alleles (risk/alternative)	Asthma (n = 243)	Control (n = 134)	P value	OR	95% CI
rs3817971 rs41355746	32361133 32364052	A/G C/T	0.949 0.951	0.840 0.877	$6.285 \times 10^{-7}$ $2.456 \times 10^{-4}$	3.509 2.703	2.090–5.889 1.562–4.679
rs41441651	32363888	C/T	0.947	0.877	$6.546 \times 10^{-4}$	2.484	1.452-4.252

Summary of GWASs of atopic moderate to severe persistent asthma at *P* value  $<1.087 \times 10^{-3}$  (= 0.05/46) in the study subjects.

CI = confidence interval; GWAS = genome-wide association study; OR = odds ratio; SNP = single nucleotide polymorphism.

quadrichotomized as follows:  $<1, \ge 1$  to  $<4, \ge 4$  to <16, and  $\ge 16$ . The 3 SNPs (rs3817971, rs41355746, and rs41441651) showed significant associations with the level and grade of PC<sub>20</sub>, as well as the presence of BHR. Association results between genetic polymorphisms and BHR are depicted in Supplemental Table S3, http://links.lww.com/MD2/A620. Differences in individual values of PC<sub>20</sub> according to each genotype are presented in Figure 2.

For BDR, no significant associations were found with the 3 SNPs after comparisons between alleles.

#### 3.5. Associations with allergic sensitization

Linear regression analysis indicated that the 3 loci did not show significant associations with the serum level of total IgE. However, analyses with Der p- and Der f-specific IgE revealed significant associations (Table 4). rs3817971 showed a significant association with serum levels of Dep p- and Dep f-specific IgE under additive, dominant, and recessive models. Further, rs41441651 and rs41335746 showed a positive association under the additive and dominant models (Supplemental Table S4, http://links.lww.com/MD2/A621). Figure 3 depicts the level of mite-specific IgE by genotypes of each SNP, and reveals significant differences among genotypic variations.

## 4. Discussion

In this study, we conducted a GWAS of asthma in 243 subjects with moderate to severe persistent asthma and allergic sensitization and 134 controls. We observed that no SNP passed the genome-wide threshold of significance; however, genetic variations of *BTNL2* harbored the top significant variant (rs3817971) and *BTNL2* was associated with atopic moderate to severe persistent asthma. We also revealed that *BTNL2* was associated with asthma-related traits such as allergic sensitization and BHR.

<i>BTNL2</i> gene, a member of the immunoglobulin superfamily, is
located at the junction of the HLA class II and class III regions on
chromosome 6p21. <sup>[24]</sup> Owing to its structural homology to the
CD80/CD86 family of costimulatory proteins, BTNL2 plays an
important role in modulating costimulatory receptors involved in
T cell activation during antigen presentation by antigen
presenting cells. <sup>[25]</sup> Human BTNL2 inhibits the proliferation
of T cells and reduces the levels of cytokines, such as interleukin-2
and interferon- $\gamma$ , which are associated with T cell activation. <sup>[26–</sup>
<sup>28]</sup> BTNL2 also induces the differentiation of regulatory T cells
from mature naïve CD4 <sup>+</sup> T cells. <sup>[29,30]</sup> To date, there have been
consistent reports on its strong genetic association with
autoimmune and inflammatory diseases. For instance, sarcoido-
sis, a systemic inflammatory granulomatous disorder character-
ized by an exaggerated cellular immune response due to increased
inflammatory activity of macrophages and CD4+ helper T
cells, <sup>[31]</sup> is representative of diseases related to <i>BTNL2</i> . <sup>[32]</sup> The
truncating mutation of rs2076530 and the concomitant loss of
the IgC domain and transmembrane helix, which disrupt the
protein's membrane localization, result in impaired T cell
downregulatory function of BTNL2. <sup>[33]</sup> Moreover, BTNL2
reportedly plays a role in the genetic susceptibility to ulcerative
colitis, which represents chronic relapsing inflammatory pheno-
types of the atypical T helper 2 (Th2) immune responses affecting
mucosal epithelial surfaces. <sup>[34,35]</sup>

Asthma is a chronic inflammatory disease of the conducting airways and involves both innate and adaptive immune mechanisms through a combination of Th1, Th2, and Th17 responses.<sup>[36]</sup> It is a heterogeneous disease with respect to its severity, presence of allergic sensitization, total or specific serum IgE levels, and prognosis as well as its difference in children and adults.<sup>[1]</sup> It is likely that the genetic architecture also differs between the phenotypic subtypes of asthma. Adoption of extreme phenotypes in our study effectively showed an association

<b>1 1 1</b>			
	ie 1	-1	

Accordiation	requite	<b>af 2</b>		with	the	-	and	dograa	of	рцр
Association	results	ors	SNPS	with	tne	presence	and	aearee	ΟΤ	внк.

		PC <sub>20</sub> ; categorical					PC <sub>20</sub> ; continuous			BHR <sup>†</sup>		
SNP	Risk allele		<1	1–4	4–16	>16	P value	β	SE	P value	aOR (95% CI)	P value
		Number of cases	79	160	197	240	<.001	-12.408	2.695	<.001	3.599 (2.089-6.200)	<.001
		Proportion	0.12	0.24	0.29	0.36						
rs41355746	С	Number of cases	80	160	198	251	.005	-9.958	2.926	.001	2.704 (1.532-4.775)	.001
		Proportion	0.12	0.23	0.29	0.36						
rs41441651	С	Number of cases	80	160	197	250	.004	-9.284	2.882	.001	2.657 (1.520-4.645)	.001
		Proportion	0.12	0.23	0.29	0.36						

Regression analyses were adjusted for age and sex.

 $\beta = \beta$  estimate; aOR = adjusted odds ratio; BHR = bronchial hyperresponsiveness; CI = confidence interval; PC<sub>20</sub> = provocative concentration of methacholine inducing a 20% decrease in forced expiratory volume in 1 s; SE = standard errors; SNP = single nucleotide polymorphism.

<sup>+</sup> BHR was defined as a decrease in forced expiratory volume in 1 s of ≥20% with inhalation of <16 mg/mL methacholine.



Figure 2. Differences in  $PC_{20}$  according to genotype groups for each SNP. The y-axis represents the value of  $PC_{20}$  in mg/mL, and the x-axis represents the 3 genotyping groups for 3 SNPs of (A) rs3817971, (B) rs41335746, and (C) rs41441651. Violin plots depict the distribution of individual data in addition to the median (thick horizontal line) and interquartile range (dotted line).  $PC_{20}$  = provocative concentration of methacholine inducing a 20% decrease in forced expiratory volume in 1; SNP = single nucleotide polymorphism. NS, P > .05; \*, P < .05; \*\*\*\*, P < .0001.

Table 4	
Association results of 3 SNPs with the level of total and mite-specific serum IgE.	

			Total IgE		D	er p specific	lgE	Der f specific IgE			
SNP	Risk allele	β	SE	P value	β	SE	P value	β	SE	P value	
rs3817971	А	170.315	91.430	.068	13.930	4.776	.004	19.075	5.960	.001	
rs41355746	С	149.837	98.680	.129	11.511	5.196	.027	14.958	6.501	.022	
rs41441651	С	142.387	97.147	.143	10.516	5.101	.040	11.957	6.388	.062	

Analyses were adjusted for age and sex.

β = β estimate; Der f = Dermatophagoides farina; Der p = Dermatophagoides pteronyssinus; IgE = immunoglobulin E; SE = standard errors; SNP = single nucleotide polymorphism.



Figure 3. Differences in mite-specific serum IgE levels according to genotyping groups for each SNP. Serum specific IgE levels towards Der p (A–C) and Der f (D–F) according to genotypes of each SNP are presented. Violin plots depict the distribution of individual data in addition to the median (thick horizontal line) and interquartile range (dotted line). IgE = immunoglobulin E; SNP = single nucleotide polymorphism; Der p = *Dermatophagoides pteronyssinus*; Der f = *Dermatophagoides farina*. NS, P > .05; \*, P < .05; \*\*, P < .01; \*\*\*

between genotypes and phenotypes. As we targeted moderate-tosevere atopic asthma in this study, BTNL2 could have been associated with asthma or allergic sensitization. All asthma cases had allergic sensitization; however, no controls had allergic sensitization. Here, we demonstrated that 3 SNPs of BTNL2 were associated with a risk of sensitization toward Der p and Der f. This finding is consistent with that of a previous report by Konno et al,<sup>[37]</sup> which suggests that *BTNL2* is a candidate gene responsible for the pathogenesis of Der f-specific IgE responsiveness. Our results also support previous observations that suggest that non-HLA genes are important for regulating specific IgEresponsiveness to complex allergens such as Der p and Der f.<sup>[38,39]</sup> Unlike other low molecular weight allergens that show particular HLA class II-associated IgE responsiveness,<sup>[40]</sup> complex allergens expressing multiple epitopes and sequence motifs<sup>[41,42]</sup> are expected to be related to more complex immune mechanisms.<sup>[43]</sup>

*BTNL2* polymorphisms were also significantly associated with the presence and degree of BHR in this study. As BHR is one of the important characteristics in asthma pathology, this novel finding also implicates the association between *BTNL2* and asthma. Indeed, there have been a few reports from other populations that may reinforce the association of *BTNL2* with asthma. Hirota et al<sup>[44]</sup> identified several susceptibility loci including rs3117098, which is located within the intron of *BTNL2*, from the GWAS of adult asthma cases in a Japanese cohort. Further, rs3117098 was found to be significantly associated with asthma susceptibility in adult Chinese Zhuang population.<sup>[45]</sup> Our 3 GWAS-supported variants, that is, these non-coding regions, may play important roles in the regulation of gene activity, thereby affecting disease development and susceptibility.<sup>[46]</sup>

The present study has several fundamental limitations. Our GWAS had a small sample size; inevitably, no significant variants were found. A GWAS using extreme phenotypes of both cases and controls cannot overcome the limitation of a small sample size.<sup>[47]</sup> However, we acknowledge that the statistical approach of using a Bonferroni-corrected P value to control the false positive rate and the stringent significance threshold would miss many true associations in a GWAS.<sup>[20]</sup> Thus, variants with small P values that do not meet genome-wide significance thresholds in GWASs remain to be further investigated as true associations. Additionally, our results should be interpreted with caution because we did not validate the identified variants in an independent cohort. Furthermore, considering BTNL2 is located on chromosome 6p21.3 near the HLA gene cluster, we cannot exclude the possibility that the identified BTNL2-asthma association was merely a reflection of LD with HLA genes. Moreover, this case-control study only provides a hypothesis of the epidemiological etiology. The potential functional mechanisms via which genetic polymorphisms may lead to asthma pathogenesis have not yet been elucidated and thus warrant further investigations.

In conclusion, we observed that *BTNL2* was associated with atopic moderate to severe persistent asthma in Korean children. *BTNL2* was also associated with asthma-related traits, including allergic sensitization to mite and BHR. To our knowledge, this is the first study in a Korean population that has demonstrated an association between the genetic variants of *BTNL2* with childhood asthma and asthma-related traits using a GWAS. However, further studies are needed to clarify the immunogenetic basis of this association with childhood asthma.

#### Author contributions

Conceptualization, Soo Yeon Kim, Min Jung Kim, Yoon Hee Kim, Myung Hyun Sohn and Kyung Won Kim; Data curation, Ga Eun Kim, Min Jung Kim, Yoon Hee Kim and Kyung Won Kim; Formal analysis, Eun Gyul Kim, Mi Na Kim and Jung Yeon Hong; Funding acquisition, Kyung Won Kim; Investigation, Soo Yeon Kim, Jae Hwa Jung and Mireu Park; Methodology, Soo Yeon Kim, Eun Gyul Kim, Ga Eun Kim, Jae Hwa Jung and Mireu Park; Supervision, Myung Hyun Sohn and Kyung Won Kim; Writing – original draft, Soo Yeon Kim; Writing – review & editing, Kyung Won Kim. All authors have read and agreed to the published version of the manuscript.

- Conceptualization: Soo Yeon Kim, Min Jung Kim, Yoon Hee Kim, Myung Hyun Sohn, Kyung Won Kim.
- Data curation: Ga Eun Kim, Min Jung Kim, Yoon Hee Kim, Kyung Won Kim.

Formal analysis: Eun Gyul Kim, Mi Na Kim, Jung Yeon Hong. Funding acquisition: Kyung Won Kim.

- Investigation: Soo Yeon Kim, Jae Hwa Jung, Mireu Park.
- Methodology: Soo Yeon Kim, Eun Gyul Kim, Ga Eun Kim, Jae Hwa Jung, Mireu Park.
- Supervision: Myung Hyun Sohn, Kyung Won Kim.
- Writing original draft: Soo Yeon Kim.
- Writing review & editing: Kyung Won Kim.

#### References

- Anderson GP. Endotyping asthma: new insights into key pathogenic mechanisms in a complex, heterogeneous disease. Lancet 2008;372:1107–19.
- [2] Moffatt MF, Gut IG, Demenais F, et al. A large-scale, consortium-based genomewide association study of asthma. N Engl J Med 2010;363: 1211–21.
- [3] Pickrell JK, Berisa T, Liu JZ, Ségurel L, Tung JY, Hinds DA. Detection and interpretation of shared genetic influences on 42 human traits. Nat Genet 2016;48:709–17.
- [4] Kim KW, Ober C. Lessons learned from GWAS of asthma. Allergy Asthma Immunol Res 2018;11:170–87.
- [5] Drake KA, Torgerson DG, Gignoux CR, et al. A genome-wide association study of bronchodilator response in Latinos implicates rare variants. J Allergy Clin Immunol 2014;133:370–8. e315.
- [6] Nieuwenhuis M, Vonk J, Himes B, et al. PTTG 1 IP and MAML 3, novel genomewide association study genes for severity of hyperresponsiveness in adult asthma. Allergy 2017;72:792–801.
- [7] Gudbjartsson DF, Bjornsdottir US, Halapi E, et al. Sequence variants affecting eosinophil numbers associate with asthma and myocardial infarction. Nat Genet 2009;41:342–7.
- [8] Weidinger S, Gieger C, Rodriguez E, et al. Genome-wide scan on total serum IgE levels identifies FCER1A as novel susceptibility locus. PLoS Genet 2008;4:e1000166.
- [9] Bønnelykke K, Matheson MC, Pers TH, et al. Meta-analysis of genomewide association studies identifies ten loci influencing allergic sensitization. Nat Genet 2013;45:902–6.
- [10] Li JZ, Absher DM, Tang H, et al. Worldwide human relationships inferred from genome-wide patterns of variation. Science 2008;319: 1100-4.
- [11] Shriner D, Adeyemo A, Gerry NP, et al. Transferability and fine-mapping of genome-wide associated loci for adult height across human populations. PLoS One 2009;4:e8398.
- [12] Baye TM, Butsch Kovacic M, Biagini Myers JM, et al. Differences in candidate gene association between European ancestry and African American asthmatic children. PLoS One 2011;6:e16522.
- [13] Kim SH, Cho BY, Park CS, et al. Alpha-T-catenin (CTNNA3) gene was identified as a risk variant for toluene diisocyanate-induced asthma by genome-wide association analysis. Clin Exp Allergy 2009;39: 203–12.
- [14] Kim J-H, Park B-L, Cheong HS, et al. Genome-wide and follow-up studies identify CEP68 gene variants associated with risk of aspirinintolerant asthma. PLoS One 2010;5:e13818.

- [15] Shin SW, Park J, Kim YJ, et al. A highly sensitive and specific genetic marker to diagnose aspirin-exacerbated respiratory disease using a genome-wide association study. DNA Cell Biol 2012;31:1604–9.
- [16] Park BL, Kim T-H, Kim J-H, et al. Genome-wide association study of aspirin-exacerbated respiratory disease in a Korean population. Human Genetics 2013;132:313–21.
- [17] Kim S-H, Cho B-Y, Choi H, et al. The SNP rs3128965 of HLA-DPB1 as a genetic marker of the AERD phenotype. PLoS One 2014;9:e111220.
- [18] Park S-Y, Jung H, Kim J-H, et al. Longitudinal analysis to better characterize Asthma-COPD overlap syndrome: findings from an adult asthma cohort in Korea (COREA). Clin Exp Allergy 2019;https://doi. org/10.1111/cea.13339.
- [19] Kim J-H, Cheong HS, Park JS, et al. A genome-wide association study of total serum and mite-specific IgEs in asthma patients. PLoS One 2013;8: e71958.
- [20] Ober C. Asthma genetics in the post-GWAS era. Ann Am Thorac Soc 2016;13(Suppl 1):S85–90.
- [21] Purcell S, Neale B, Todd-Brown K, et al. PLINK: a tool set for wholegenome association and population-based linkage analyses. Am J Hum Genet 2007;81:559–75.
- [22] Turner SD. qqman: an R package for visualizing GWAS results using QQ and manhattan plots. Biorxiv 2014;5165.
- [23] Pruim RJ, Welch RP, Sanna S, et al. LocusZoom: regional visualization of genome-wide association scan results. Bioinformatics 2010;26: 2336–7.
- [24] Traherne JA, Barcellos LF, Sawcer SJ, et al. Association of the truncating splice site mutation in BTNL2 with multiple sclerosis is secondary to HLA-DRB1\*15. Hum Mol Genet 2006;15:155–61.
- [25] Arnett HA, Escobar SS, Viney JL. Regulation of costimulation in the era of butyrophilins. Cytokine 2009;46:370–5.
- [26] Nguyen T, Liu XK, Zhang Y, Dong C. BTNL2, a butyrophilin-like molecule that functions to inhibit T cell activation. J Immunol 2006; 176:7354–60.
- [27] Arnett HA, Escobar SS, Gonzalez-Suarez E, et al. BTNL2, a butyrophilin/ B7-like molecule, is a negative costimulatory molecule modulated in intestinal inflammation. J Immunol 2007;178:1523–33.
- [28] Yamazaki T, Goya I, Graf D, Craig S, Martin-Orozco N, Dong C. A butyrophilin family member critically inhibits T cell activation. J Immunol 2010;185:5907–14.
- [29] Ammann JU, Cooke A, Trowsdale J. Butyrophilin Btn2a2 inhibits TCR activation and phosphatidylinositol 3-kinase/Akt pathway signaling and induces Foxp3 expression in T lymphocytes. J Immunol 2013;190: 5030–6.
- [30] Swanson RM, Gavin MA, Escobar SS, et al. Butyrophilin-like 2 modulates B7 costimulation to induce Foxp3 expression and regulatory T cell development in mature T cells. J Immunol 2013;190:2027–35.
- [31] Newman LS, Rose CS, Maier LA. Medical progress: sarcoidosis. N Engl J Med 1997;336:1224–34.

- [32] Calender A, Weichhart T, Valeyre D, Pacheco Y. Current insights in genetics of sarcoidosis: functional and clinical impacts. J Clin Med 2020;9:2633.
- [33] Valentonyte R, Hampe J, Huse K, et al. Sarcoidosis is associated with a truncating splice site mutation in BTNL2. Nat Genet 2005; 37:357-64.
- [34] Mochida A, Kinouchi Y, Negoro K, et al. Butyrophilin-like 2 gene is associated with ulcerative colitis in the Japanese under strong linkage disequilibrium with HLA-DRB1\* 1502. Tissue Antig 2007;70:128–35.
- [35] Pathan S, Gowdy R, Cooney R, et al. Confirmation of the novel association at the BTNL2 locus with ulcerative colitis. Tissue Antig 2009;74:322–9.
- [36] Lambrecht BN, Hammad H. The immunology of asthma. Nat Immunol 2015;16:45–56.
- [37] Konno S, Takahashi D, Hizawa N, et al. Genetic impact of a butyrophilin-like 2 (BTNL2) gene variation on specific IgE responsiveness to Dermatophagoides farinae (Der f) in Japanese. Allergol Int 2009;58:29–35.
- [38] Hizawa N, Freidhoff LR, Ehrlich E, et al. Genetic regulation of Dermatophagoides pteronyssinus-specific IgE responsiveness: a genomewide multipoint linkage analysis in families recruited through 2 asthmatic sibs. J Allergy Clin Immunol 1998;102:436–42.
- [39] Blumenthal M, Ober C, Beaty T, et al. Genome scan for loci linked to mite sensitivity: the Collaborative Study on the Genetics of Asthma (CSGA). Genes Immun 2004;5:226–31.
- [40] Huang SK, Zwollo P, Marsh DG. Class II major histocompatibility complex restriction of human T cell responses to short ragweed allergen, Amb a V. Eur J Immunol 1991;21:1469–73.
- [41] Yssel H, Johnson K, Schneider P, et al. T cell activation-inducing epitopes of the house dust mite allergen Der p I. Proliferation and lymphokine production patterns by Der p I-specific CD4+ T cell clones. J Immunol 1992;148:738–45.
- [42] van Neerven RJ, van t'Hof W, Ringrose J, et al. T cell epitopes of house dust mite major allergen Der p II. J Immunol 1993;151:2326–35.
- [43] Marsh DG, Bias WB, Ishizaka K. Genetic control of basal serum immunoglobulin E level and its effect on specific reaginic sensitivity. Proc Natl Acad Sci 1974;71:3588–92.
- [44] Hirota T, Takahashi A, Kubo M, et al. Genome-wide association study identifies three new susceptibility loci for adult asthma in the Japanese population. Nat Genet 2011;43:893–6.
- [45] Liang SQ, Deng JM, Wei X, et al. Association of GWAS-supported noncoding area loci rs404860, rs3117098, and rs7775228 with asthma in Chinese Zhuang population. J Clin Lab Anal 2020;34:e23066.
- [46] Gloss BS, Dinger ME. Realizing the significance of noncoding functionality in clinical genomics. Exp Mol Med 2018;50:1–8.
- [47] Kim KW, Myers RA, Lee JH, et al. Genome-wide association study of recalcitrant atopic dermatitis in Korean children. J Allergy Clin Immunol 2015;136:678–84. e674.